

Supplemental Material

Manuscript title: Advancing the Next Generation of Risk Assessment

Ila Cote,¹ Melvin Andersen,² Gerald Ankley,¹ Stanley Barone,¹ Linda Birnbaum,³ Kim Boekelheide,⁴ Frederic Bois,⁵ Lyle Burgoon,¹ Weihsueh A. Chiu,⁶ Douglas Crawford-Brown,⁷ Kevin M. Crofton,¹ Michael DeVito,³ Robert Devlin,¹ Stephen Edwards,¹ Kathryn Z. Guyton,⁸ Dale Hattis,⁹ Richard Judson,¹ Derek Knight,¹⁰ Daniel Krewski,¹¹ Jason Lambert,¹ Elizabeth Maull,³ Donna Mendrick,¹² Gregory M. Paoli,¹³ Chirag Patel,¹⁴ Edward Perkins,¹⁵ Gerald Poje,¹⁶ Christopher Portier,¹⁷ Ivan Rusyn,⁶ Paul Schulte,¹⁸ Anton Simeonov,¹⁹ Martyn Smith,²⁰ Kristina A. Thayer,³ Russell S. Thomas,¹ Reuben Thomas,²⁰ Raymond R. Tice,³ John Vandenberg,¹ Daniel L. Villeneuve,¹ Scott Wesselkamper,¹ Maurice Whelan,²¹ Christine Whittaker,¹⁸ Ronald H. White²² Menghang Xia,¹⁹ Carole Yauk,²³ Lauren Zeise,²⁴ Q. Jay Zhao,¹ Robert S. DeWoskin¹

¹U.S. Environmental Protection Agency

²The Hamner Institutes for Health Sciences

³National Institute of Environmental Health Sciences and the National Toxicology Program

⁴Brown University

⁵L'Institut National de l'Environnement Industriel et des Risques

⁶Texas A & M University

⁷University of Cambridge, Department of Land Economy,

⁸International Agency for Cancer Research

⁹Clark University

¹⁰European Chemicals Agency

¹¹University of Ottawa

¹²Food and Drug Administration, National Center for Toxicological Research

¹³Risk Sciences International

¹⁴Harvard University

¹⁵Army Corps of Engineers

¹⁶Grant Consulting Group

¹⁷Environmental Defense Fund

¹⁸National Institute for Occupational Safety and Health

¹⁹National Institutes of Health, National Center for Advancing Translational Science

²⁰University of California, Berkeley

²¹European Commission Joint Research Centre, Institute for Health and Consumer Protection

²²Center for Effective Government

²³Health Canada

²⁴California Environmental Protection Agency, Office of Environmental Health Hazard Assessment

Table of Contents

Table S1. Governmental Partners	3
Table S2: An Illustrative Framework for Evidence Integration Focusing on New Data Types Presented in the NexGen Report.	4

Table S1. Governmental Partners

- Army Corps of Engineers
- California Environmental Protection Agency, Office of Environmental Health Hazard Assessment
- European Chemicals Agency
- European Joint Research Centre
- Health Canada
- L'Institut National de l'Environnement Industriel et des Risques
- U.S. Food and Drug Administration (National Center for Toxicological Research)
- U.S. Centers for Disease Control and Prevention, National Center for Environmental Health, and Agency for Toxic Substances and Disease Registry
- U.S. Department of Defense
- U.S. National Center for Advancing Translational Science
- U.S. National Institute of Environmental Health Sciences and National Toxicology Program
- U.S. National Institute for Occupational Safety and Health

Table S2: An Illustrative Framework for Evidence Integration Focusing on New Data Types Presented in the NexGen Report.

This causal determination framework illustrates how evidence integration and inferences about causality could be made using new data types. The left column summarizes the prototype results, the middle column presents evidence for causality exemplified by the prototypes, and the right column illustrates how such prototypic evidence might be integrated and weighed. The first set of prototypes is unique in that the prototypes have known human health effects and well-documented public health risks. For these prototypes, the “Evidence Integration” column evaluates how successful new data types were in predicting known outcomes (criteria for study selection, evidence integration and causal determination considered here are discussed in DHHS 2014; EPA 2013a; McConnell et al. 2014; Meek et al. 2014; NRC 2014; Rhomberg et al. 2013). Modifications of the Bradford-Hill criteria (e.g., consistency, coherence, biologic plausibility) continued to be useful in the evaluation of new data types. As presented here, confidence in causality ranges from suggestive to likely, largely based on the understanding of the biologic context in which new data types are embedded. “Likely” is generally for the limited number of cases where the new data types are well anchored to adverse outcomes by a combination of observational and experimental data, and include mechanistic and systems biology understanding. Biologic context need not be chemical specific but can be derived from disease-/disorder-specific knowledge or from analogy with related chemicals. In practice, much of the new datatypes are anticipated to be suggestive for the near-term and most appropriate for screening and prioritization and, perhaps, limited-scope assessments. Of note is that, contrary to traditional approaches, some new approaches can be used to estimate relative potencies or toxicity values in the absence of clearly identified hazards. Major assessments are anticipated to be augmented by new data types but, for the near term, continue to be based on traditional data. To simplify the table, similar prototypes with shared attributes are aggregated where possible.

Table S2: An Illustrative Framework for Evidence Integration Focusing on New Data Types Presented in the NexGen Report.		
Prototypes	Evidence for Causality	Evidence Integration

Tie r 3	<p>Molecular epidemiology and clinical studies:</p> <ul style="list-style-type: none"> • Illustrated that new data types (when properly collected, analyzed, and reported) appear to provide results comparable to robust, traditional human data, and could be used, when linked to mechanistic information, to: (1) evaluate potential hazard posed by chemicals with no or limited traditional data, (2) augment traditional assessments, or (3) better inform traditional risk assessment issues, such as human variability and susceptibility, cross-species and <i>in vivo</i>-to-<i>in vitro</i> comparisons, cumulative risk, and low exposure-dose-response relationships. 	<ul style="list-style-type: none"> • Specific pattern alterations in molecular events appear consistently and strongly associated with known intermediate events and known hazards at environmental exposure levels (EPA 2013b, 2015a; McCullough et al. 2014; McHale et al. 2009; McHale et al. 2012; Smith et al. 2011; Thomas et al. 2014). Data for tobacco smoke were reported in the Gene Expression Omnibus (GEO) or ArrayExpress (AE). Also see Cooper et al. (2013). • Exposure-dose was measured for benzene and ozone using urinary biomarkers and radiolabeled ozone, respectively (Hatch et al. 2014; McHale et al. 2009; Thomas et al. 2014). Tobacco-smoke exposures were self-reported, substantially increasing uncertainty for exposure-dose-response characterization and highlighting the need for accurate exposure characterization (EPA 2015a). • Dose-dependent alterations are observed in concomitantly collected molecular events and adverse effects, in the range of environmental exposure (benzene and ozone). Some molecular pathways are altered at all concentrations; other molecular and toxicological effects emerge with increasing dose. Molecular patterns that occur consistently across all concentrations appear preferable as biomarkers (EPA 2013b; Hatch et al. 2014; McCullough et al. 2014; Thomas et al. 2014). • Pharmacological interventions have been shown to modify identified AOPs, and, concomitantly, the incidence or severity of the adverse outcomes (Cooper et al. 2013; Hatzimichael and Crook 2013; McCullough et al. 2014). 	<p>Suggestive to likely: Evidence is consistent, coherent, and biologically plausible that the observed molecular events are causally related to adverse effects. Implications are based on comparisons to robust traditional risk assessments:</p> <ul style="list-style-type: none"> • For benzene and ozone, identified molecular events are likely causally related to known adverse outcomes in a dose-dependent fashion. Mechanistic links between molecular events, intermediate effects, and adverse outcomes are well understood. Pharmacologic intervention that blocks implicated pathways also blocks or ameliorates adverse effects. • In comparison, the molecular data for PAH are considered suggestive of a causal association between PAH and lung cancer due to a lack of an observed exposure-dose-response relationship (likely due to uncertainties in exposure characterization), and data quality, analysis, and reporting limitations. Only about 8% of studies in GEO and AE met study selection criteria.
------------	---	---	---

		<ul style="list-style-type: none"> • Additional evidence for the involvement of specific pathways in disease is provided by identification of naturally occurring human gene variants in the AOP network that alter susceptibility and risks (Cooper et al. 2013; Hatzimichael and Crook 2013; Moreno-Macias et al. 2013; Schlenk et al. 2008; Shen et al. 2011; Sillé et al. 2012; Smith et al. 2011; Vawda et al. 2014; Zhuo et al. 2012). • Adverse outcome pathway (AOP) networks are also disrupted by other chemical and nonchemical stressors known to alter the incidence of the specific disease/disorder under consideration; thus, AOP networks provide a tool for evaluating cumulative risks based on mechanistic commonalities (IARC 2012; Smith et al. 2011; R Thomas et al. 2012). • Supporting data are provided by multiple molecular epidemiology and clinical studies and chronic animal bioassays and coherence with other systems biology data (NIH BioSystems: acute myeloid leukemia, lung cancer (small cell, non-small cell); Comparative Toxicogenomics Database: PAHs and cancer: (EPA 2013a, b); BaP and cancer, ozone and respiratory disease). • Although species and <i>in vitro</i> differences exist, these examples provide consistent, coherent biologically plausible data linking specific omic alterations with specific diseases. 	<p>Suggestive vs. likely: In general, molecular data alone associated with adverse outcomes are expected to be only suggestive or inadequate for causal determination. To rise to likely, the following are generally necessary: multiple, consistent, high-quality observational studies with similar results; understanding of the cascade of events between molecular events to adverse outcomes, and experimental evidence showing that reversal of pathway alterations blocks or ameliorates adverse outcome; or naturally occurring experiments where gene variants alter incidence or characteristic of disease. Important variables such as experimental paradigm (e.g., <i>in vivo</i> vs. <i>in vitro</i>), cell type, tissue type, and species also require consideration. Suggestive data are likely to be most useful for hypothesis generation, discovery, screening and prioritization, and potential augmentation of traditional data.</p>
Tier 2	<p>Knowledge mining and meta-analysis prototype illustrated how large searchable databases can be used to identify, organize, integrate, and analyze existing data in an automated (computerized) fashion to discover new insights into public health risks.</p>	<ul style="list-style-type: none"> • Knowledge mining and meta-analysis discovered associations between known exposures (biomonitoring) to several environmental agents (e.g., metals and persistent organics chemical) with prediabetes/diabetes using the Centers for Disease Control and Prevention’s National Health Assessment Examination Survey. Human tissue biomarker and clinical outcome data are from the same individuals (EPA 2015a; Patel et al. 2012; Patel et al. 2013a). • Supporting data are found elsewhere in the literature (NIEHS 2015b; Thayer et al. 2012). • No systems biology context or AOP is available for data from NHANES. Information on biological context is available from the literature but currently not easily accessible in high- or medium-throughput approaches (Audouze et al. 2013; Inadera 2013). • Prototype also explored possible links among site-specific chemical exposures, ethnicity, genetic variants, and diabetes risks (Dimas et al. 2014; EPA 2015a; Patel and Cullen 2012). 	<p>Suggestive: Could rise to likely with the types of supporting data noted above under “Suggestive vs. likely.”</p> <p>Associative data generally most useful for hypothesis generation.</p>

<p>Short-duration <i>in vivo</i> exposure bioassays data use in either alternative or rodents species is illustrated in two prototypes.</p>	<p>Short-duration, <i>in vivo</i> exposure bioassays—alternative (nonmammalian) species prototype included example zebrafish developmental assay results, characterization of thyroid-specific mechanisms, and predictive modeling of complex dose-response phenomena. Padilla et al. (2012) reported AC₅₀s in a zebrafish developmental assay for 305 chemicals. Potencies for individual chemicals and chemical classes were shown to range over several orders of magnitude (1 nM–80 µM). For certain classes of chemicals, 80–100% of the chemicals in a class tested positive (embryo death or structural defects). Perkins et al. (2013) illustrated the use of alternative species to help articulate mechanisms showing an AOP network for thyroid disruption with example toxicants and alternative models applicable to both human and ecological hazard assessment. Also discussed are how predictive models coupled to mechanistic understanding can be used to better characterize dose-response (Sipes et al. 2011b), circadian variations (Eisenberg et al. 2008), and exposure window-response relationships (DeWoskin et al. 2014). For more details on alternative species bioassays see Ankley and Gray (2013), Perkins et al. (2013), and Villeneuve et al. (2014).</p>	<p>Suggestive to likely for consistent, coherent, biologically plausible adverse phenotypic outcome data from nonmammalian, vertebrate species. Confidence is generally higher for evolutionarily conserved processes. Suggestive: For transcriptomic changes correlated to adverse outcomes studies and coupled to AOPs.</p> <p>High-content assays with measurable adverse outcomes (e.g., zebrafish developmental assay) generally have greater evidentiary weight than initiating event assays (e.g., transcriptomic assays). Some systems biology context is needed for limited-scope assessments for human risk (e.g., cross-species conservation, AOPs). Alternative species outcome data alone are sufficient for ecologic risk assessment. Cross-species extrapolation and subchronic measurement of indicators introduces additional uncertainties as compared to human data discussed above data. Reverse toxicokinetic models are needed to estimate equivalent human doses.</p>
	<p>Short-duration, <i>in vivo</i> exposure bioassays—rodent prototype correlated transcriptomic alterations with adverse outcomes, as determined in traditional bioassays for 10 chemicals (Thomas et al. 2011; RS Thomas et al. 2012; Thomas et al. 2013c). Consistency of the correlation between transcriptional changes and adverse effects across different exposure periods was also demonstrated (5 days to 13 weeks) (Thomas et al. 2013c). Transcriptional changes appeared at somewhat lower concentrations than traditional effects. Transcriptomic studies alone cannot predict specific hazards but might be useful to relatively ranking chemical potencies to induce biologic alterations that might precede adverse outcomes. In general, transcriptomic data require some biologic context (e.g., AOPs) to increase confidence of biologic significance.</p>	

Tier 1	<p>QSAR and molecular docking models are used to generate potency estimates and, with less confidence, hazards. Read-across is also considered (i.e., filling data gaps for data-poor chemicals by analogy with structurally related, more data-rich chemicals).</p>	<p>QSAR models can predict chemical-specific toxicity values based on chemical inherent properties for several data-poor chemicals.</p> <ul style="list-style-type: none"> • Models are developed based on chemical structures and known outcomes for data-rich chemicals. • Organisation for Economic Co-operation and Development (OECD) is harmonizing international use of QSAR hazard models and read-across in the OECD QSAR toolbox (OECD 2014, 2015a, b). • Often, the consensus of a suite of appropriate models is the preferred approach. • Often, models better predict potency than specific effects. • Issues exist around characterizing the uncertainty in QSAR and related read-across approaches and in the transparency of some models [see Ball et al. (2014) and Patlewicz et al. (2013)]. 	<p>Suggestive to Likely: TOPKAT Model predictions of potency when model is appropriate for chemicals evaluated; not generally predictive of dose-response for specific hazards; does generate a lowest-observed-adverse-effect level for a subset of the data-poor chemicals that meet confidence criteria. Additional OECD models and read-across can improve confidence in hazard characterization.</p>
	<p>High-throughput, <i>in vitro</i> bioassays and virtual tissue models are discussed.</p>	<p>High-throughput <i>in vitro</i> assays based on biological process disruptions are interpreted in systems biology and AOP contexts, and associated with adverse outcomes (Attene-Ramos et al. 2013; EPA 2015a; Judson et al. 2014; Tice et al. 2013). Virtual tissue modeling provides additional tools for evaluating data-limited chemicals (Knudsen et al. 2013).</p>	<p>Suggestive: When coupled with understanding of the AOP(s). Could rise to likely with the types of supporting data noted above under “Suggestive vs. likely.”</p>

References

- Ankley GT, Gray LE. 2013. Cross-species conservation of endocrine pathways: a critical analysis of tier 1 fish and rat screening assays with 12 model chemicals. *Environ Toxicol Chem* 32:1084–1087.
- Attene-Ramos MS, Miller N, Huang R, Michael S, Itkin M, Kavlock RJ, et al. 2013. The Tox21 robotic platform for the assessment of environmental chemicals - from vision to reality. *Drug Discov Today* 18:716–273.
- Audouze K, Brunak S, Grandjean P. 2013. A computational approach to chemical etiologies of diabetes. *Sci Rep* 3:2712; doi:10.1038/srep02712.
- Ball N, Bartels M, Budinsky R, Klapacz J, Hays S, Kirman C, et al. 2014. The challenge of using read-across within the EU REACH regulatory framework; how much uncertainty is too much? Dipropylene glycol methyl ether acetate, an exemplary case study. *Regul Toxicol Pharmacol* 68:212–221.
- Cooper WA, Lam DC, O'Toole SA, Minna JD. 2013. Molecular biology of lung cancer. *J Thorac Dis* 5 Suppl 5:S479–490.
- DeWoskin RS, Knudsen TB, Shah I. 2014. Virtual models (vM). In: *Encyclopedia of Toxicology*, 3rd Edition, Vol. 4 (Wexler P, ed). London, UK:Elsevier, Inc., 948–957.
- DHHS (U.S. Department of Health and Human Services). 2014. The Health Consequences of Smoking - 50 Years of Progress. Rockville, MD:Office of the Surgeon General. Available: <http://www.surgeongeneral.gov/library/reports/50-years-of-progress/exec-summary.pdf> [accessed 28 April 2015].
- Dimas AS, Lagou V, Barker A, Knowles JW, Magi R, Hivert MF, et al. 2014. Impact of type 2 diabetes susceptibility variants on quantitative glycemic traits reveals mechanistic heterogeneity. *Diabetes* 63:2158–2171.
- Eisenberg MC, Samuels M, DiStefano JJ, 3rd. 2008. Extensions, validation, and clinical applications of a feedback control system simulator of the hypothalamo-pituitary-thyroid axis. *Thyroid* 18:1071–1085.
- EPA (U.S. Environmental Protection Agency). 2013a. IRIS Toxicological Review of Benzo[a]pyrene (Public Comment Draft). Washington, DC:EPA. Available: http://cfpub.epa.gov/ncea/iris_drafts/recorddisplay.cfm?deid=66193 [accessed 28 April 2015].
- EPA (U.S. Environmental Protection Agency). 2013b. Integrated Science Assessment of Ozone and Related Photochemical Oxidants (Final Report). EPA/600/R-10/076F. Washington, DC:U.S. EPA. Available: <http://cfpub.epa.gov/ncea/isa/recorddisplay.cfm?deid=247492> [accessed 28 April 2015].
- EPA (U.S. Environmental Protection Agency). 2015a. Basic Information about the Toxic Substance Control Act Inventory. Available: <http://www.epa.gov/oppt/existingchemicals/pubs/tscainventory/basic.html#background> [accessed 28 April 2015].
- Hatch GE, Duncan KE, Diaz-Sanchez D, Schmitt MT, Ghio AJ, Carraway MS, et al. 2014. Progress in assessing air pollutant risks from *in vitro* exposures: matching ozone dose and effect in human airway cells. *Toxicol Sci* 141:198–205.
- Hatzimichael E, Crook T. 2013. Cancer epigenetics: New therapies and new challenges. *J Drug Deliv* 2013:529312;

doi:10.1155/2013/529312:529312.

IARC. 2012. Benzene. IARC Monogr Eval Carcinog Risk Hum 100F:249–294.

Inadera H. 2013. Developmental origins of obesity and type 2 diabetes: molecular aspects and role of chemicals. *Environ Health Prev Med* 18:185–197.

Judson R, Houck K, Martin M, Knudsen TB, Thomas RS, Sipes N, et al. 2014. *In vitro* and modelling approaches to risk assessment from the U.S. Environmental Protection Agency ToxCast programme. *Basic Clin Pharmacol Toxicol* 115:69–76.

Knudsen T, Martin M, Chandler K, Kleinstreuer N, Judson R, Sipes N. 2013. Predictive models and computational toxicology. *Methods Mol Biol* 947:343–374.

McConnell ER, Bell SM, Cote I, Wang RL, Perkins EJ, Garcia-Reyero N, et al. 2014. Systematic Omics Analysis Review (SOAR) tool to support risk assessment. *PloS One* 9:e110379; doi:10.1371/journal.pone.0110379.

McCullough SD, Duncan KE, Swanton SM, Dailey LA, Diaz-Sanchez D, Devlin RB. 2014. Ozone induces a pro-inflammatory response in primary human bronchial epithelial cells through mitogen-activated protein kinase activation without nuclear factor- κ B activation. *Am J Respir Cell Mol Biol* 51:426–435.

McHale CM, Zhang L, Lan Q, Li G, Hubbard AE, Forrest MS, et al. 2009. Changes in the peripheral blood transcriptome associated with occupational benzene exposure identified by cross-comparison on two microarray platforms. *Genomics* 93:343–349.

McHale CM, Zhang L, Smith MT. 2012. Current understanding of the mechanism of benzene-induced leukemia in humans: Implications for risk assessment. *Carcinogenesis* 33:240–252.

Meek ME, Boobis A, Cote I, Dellarco V, Fotakis G, Munn S, et al. 2014. New developments in the evolution and application of the WHO/IPCS framework on mode of action/species concordance analysis. *J Appl Toxicol* 34:1–18.

Moreno-Macias H, Dockery DW, Schwartz J, Gold DR, Laird NM, Sienra-Monge JJ, et al. 2013. Ozone exposure, vitamin C intake, and genetic susceptibility of asthmatic children in Mexico City: a cohort study. *Respir Res* 14:14; doi:10.1186/1465-9921-14-14.

NIEHS (National Institute of Environmental Health Sciences). 2015b. Comparative Toxicogenomic Database (CTD)TM. Available: <http://ctdbase.org/> [accessed 28 April 2015].

NRC (National Research Council). 2014. Review of EPA's Integrated Risk Information System (IRIS) Process. Washington, DC: The National Academies Press. Available: http://www.nap.edu/catalog.php?record_id=18764 [accessed 28 April 2015].

OECD (Organisation for Economic Cooperation and Development). 2014. The OECD QSAR Toolbox. Version 3.0. Available: <http://www.oecd.org/env/chemicalsafetyandbiosafety/assessmentofchemicals/theoecdqsartoolbox.htm> [accessed 29 August 2014].

OECD (Organisation for Economic Cooperation and Development). 2015a. Other Activities on Molecular Screening and Toxicogenomics. Available: <http://www.oecd.org/env/ehs/testing/toxicogenomics.htm> [accessed 28 April 2015].

OECD (Organisation for Economic Cooperation and Development). 2015b. Grouping of Chemicals: Chemical Categories and Read-Across. Available: <http://www.oecd.org/chemicalsafety/risk-assessment/groupingofchemicalschemicalcategoriesandread-across.htm> [accessed 28 April 2015].

Padilla S, Corum D, Padnos B, Hunter DL, Beam A, Houck KA, et al. 2012. Zebrafish developmental screening of the ToxCastTM

Phase I chemical library. *Reprod Toxicol* 33:174–187.

Patel CJ, Chen R, Butte AJ. 2012. Data-driven integration of epidemiological and toxicological data to select candidate interacting genes and environmental factors in association with disease. *Bioinformatics* 28:i121–126.

Patel CJ, Cullen MR. 2012. Genetic variability in molecular responses to chemical exposure. *EXS* 101:437–457.

Patel CJ, Chen R, Kodama K, Loannidis JP, Butte AJ. 2013a. Systematic identification of interaction effects between genome-and environment-wide associations in type 2 diabetes mellitus. *Hum Genet* 132:495–508.

Patlewicz G, Ball N, Booth ED, Hulzebos E, Zvinavashe E, Hennes C. 2013. Use of category approaches, read-across and (Q)SAR: general considerations. *Regul Toxicol Pharmacol* 67:1–12.

Perkins EJ, Ankley GT, Crofton KM, Garcia-Reyero N, Lalone CA, Johnson MS, et al. 2013. Current perspectives on the use of alternative species in human health and ecological risk assessments. *Environ Health Perspect* 121:1002–1010.

Rhomberg LR, Goodman JE, Bailey LA, Prueitt RL, Beck NB, Bevan C, et al. 2013. A survey of frameworks for best practices in weight-of-evidence analyses. *Crit Rev Toxicol* 43:753–784.

Schlenk RF, Dohner K, Krauter J, Frohling S, Corbacioglu A, Bullinger L, et al. 2008. Mutations and treatment outcome in cytogenetically normal acute myeloid leukemia. *N Engl J Med* 358:1909–1918.

Shen M, Zhang L, Lee KM, Vermeulen R, Hosgood HD, Li G, et al. 2011. Polymorphisms in genes involved in innate immunity and susceptibility to benzene-induced hematotoxicity. *Exp Mol Med* 43:374–378.

Sillé FC, Thomas R, Smith MT, Conde L, Skibola CF. 2012. Post-GWAS functional characterization of susceptibility variants for chronic lymphocytic leukemia. *PloS One* 7:e29632; doi:10.1371/journal.pone.0029632:e29632.

Sipes NS, Padilla S, Knudsen TB. 2011b. Zebrafish: as an integrative model for twenty-first century toxicity testing. *Birth Defects Res C Embryo Today* 93:256–267.

Smith MT, Zhang L, McHale CM, Skibola CF, Rappaport SM. 2011. Benzene, the exposome and future investigations of leukemia etiology. *Chem Biol Interact* 192:155–159.

Thayer KA, Heindel JJ, Bucher JR, Gallo MA. 2012. Role of environmental chemicals in diabetes and obesity: a National Toxicology Program workshop review. *Environ Health Perspect* 120:779–789.

Thomas R, Phuong J, McHale CM, Zhang L. 2012. Using bioinformatic approaches to identify pathways targeted by human leukemogens. *Int J Environ Res Public Health* 9:2479–2503.

Thomas R, Hubbard A, McHale CM, Zhang L, Rappaport SM, Rothman N, et al. 2014. Characterization of changes in gene expression and biochemical pathways at low levels of benzene exposure. *PloS One* 9:e91828; doi:10.1371/journal.pone.0091828.

Thomas RS, Clewell HJ, 3rd, Allen BC, Wesselkamper SC, Wang NC, Lambert JC, et al. 2011. Application of transcriptional benchmark dose values in quantitative cancer and noncancer risk assessment. *Toxicol Sci* 120:194–205.

Thomas RS, Clewell HJ, 3rd, Allen BC, Yang L, Healy E, Andersen ME. 2012. Integrating pathway-based transcriptomic data into quantitative chemical risk assessment: a five chemical case study. *Mutat Res* 746:135–143.

Thomas RS, Wesselkamper SC, Wang NC, Zhao QJ, Petersen DD, Lambert JC, et al. 2013. Temporal concordance between apical

and transcriptional points of departure for chemical risk assessment. *Toxicol Sci* 134:180–194.

Tice RR, Austin CP, Kavlock RJ, Bucher JR. 2013. Improving the human hazard characterization of chemicals: a Tox21 update. *Environ Health Perspect* 121:756–765.

Vawda S, Mansour R, Takeda A, Funnell P, Kerry S, Mudway I, et al. 2014. Associations between inflammatory and immune response genes and adverse respiratory outcomes following exposure to outdoor air pollution: a HuGE systematic review. *Am J Epidemiol* 179:432–442.

Villeneuve D, Volz DC, Embry MR, Ankley GT, Belanger SE, Leonard M, et al. 2014. Investigating alternatives to the fish early-life stage test: a strategy for discovering and annotating adverse outcome pathways for early fish development. *Environ Toxicol Chem* 33:158–169.

Zhuo W, Zhang L, Zhu B, Qiu Z, Chen Z. 2012. Association between CYP1A1 Ile462Val variation and acute leukemia risk: meta-analyses including 2164 cases and 4160 controls. *PloS One* 7:e46974; doi:10.1371/journal.pone.0046974.